arXiv:0812.0020v1 [math.FA] 28 Nov 2008

# Trace Operators for Modulation, $\alpha$-Modulation and Besov Spaces

**Hans G. Feichtinger**$^{a,1}$, **Chunyan Huang**$^{b,2}$, **Baoxiang Wang**$^{b,3}$

$^{a}$*Faculty of Mathematics, University of Vienna, Vienna A-1090, Austria .*
$^{b}$*LMAM, School of Mathematical Sciences, Peking University, Beijing 100871, China.*

**Abstract.** In this paper, we consider the trace theorem for modulation spaces $M^s_{p,q}$, $\alpha$-modulation spaces $M^{s,\alpha}_{p,q}$ and Besov spaces $B^s_{p,q}$. For the modulation space, we obtain the sharp results.

**Key words and phrases.** Trace theorems; modulation spaces; $\alpha$-modulation spaces; Besov spaces.

**2000 Mathematics Subject Classifications.** 42B35, 46E35.

## 1 Introduction

The $\alpha$-modulation spaces $M^{s,\alpha}_{p,q}$, introduced by Gröbner in [10] are a class of function spaces that contain Besov spaces $B^s_{p,q}$ ($\alpha = 1$) and modulation spaces $M^s_{p,q}$ ($\alpha = 0$) as special cases.

There are two kinds of basic coverings on Euclidean $\mathbb{R}^n$ which is very useful in the theory of function spaces and their applications, one is the uniform covering $\mathbb{R}^n = \bigcup_{k\in\mathbb{Z}^n} Q_k$, where $Q_k$ denote the unit cube with center $k$; another is the dyadic covering $\mathbb{R}^n = \bigcup_{k\in\mathbb{N}}\{\xi : 2^{k-1} \leqslant |\xi| < 2^k\}\bigcup\{\xi : |\xi| \leqslant 1\}$. Roughly speaking, these decompositions together with the frequency-localized techniques yield the frequency-uniform decomposition operator $\Box_k \sim \mathscr{F}^{-1}\chi_{Q_k}\mathscr{F}$ and the dyadic decomposition operator $\Delta_k \sim \mathscr{F}^{-1}\chi_{\{\xi:|\xi|\sim 2^k\}}\mathscr{F}$, respectively. The tempered distributions acted on these decomposition operators and equipped with the $\ell^q(L^p(\mathbb{R}^n))$ norms, we then obtain Feichtinger's modulation spaces and Besov spaces, respectively.

During the past twenty years, the third covering was independently found by Feichtinger and Gröbner [3, 4, 10], and Päivärinta and Somersalo [12]. This covering, so called $\alpha$-covering has a moderate scale which is rougher than that of the uniform covering and is thinner than that of the dyadic covering. Applying the $\alpha$-covering

---

[1] Email: hans.feichtinger@univie.ac.at
[2] Email: chunyanh80@gmail.com
[3] Email: wbx@math.pku.edu.cn

to the frequency spaces, in a similar way as the definition of Besov spaces, Gröbner [10] introduced the notion of $\alpha$-modulation spaces.

Let $n \geqslant 2$. For any $x = (x_1, ..., x_n) \in \mathbb{R}^n$, we denote $\bar{x} = (x_1, ..., x_{n-1})$. Given a Banach function space $X(\mathbb{R}^n)$ defined on $\mathbb{R}^n$ and $f \in X$, we ask for the trace of $f$ on the hyperplane $\{x : x = (\bar{x}, 0)\}$. For the sake of convenience, this hyperplane will be written as $\mathbb{R}^{n-1}$. It is clear that a clarification of this problem is of importance for the boundary value problems of the partial differential equations. Now we exactly describe the trace operators.

**Definition 1.1** *Let $X$ and $Y$ be quasi-Banach function spaces defined on $\mathbb{R}^n$ and $\mathbb{R}^{n-1}$, respectively. Denote*

$$\mathbb{Tr} : \; f(x) \to f(\bar{x}, 0). \tag{1.1}$$

*If $\mathbb{Tr} : X \to Y$ and there exists a constant $C > 0$ such that*

$$\|\mathbb{Tr} f(x)\|_Y \leqslant C\|f\|_X, \quad \forall f \in X, \tag{1.2}$$

*and there exist a continuous linear operator $\mathbb{Tr}^{-1} : Y \to X$ such that $\mathbb{Tr}\mathbb{Tr}^{-1}$ is identical operator, then $\mathbb{Tr}$ is said to be a retraction from $X$ onto $Y$.*

If $\mathbb{Tr}$ is a retraction from $Y$ onto $X$, we see that the trace of $f \in X$ is well behaved in $Y$. The trace theorems in modulation spaces and Besov spaces have been extensively studied. Feichtinger [5] considered the trace theorem for the modulation space $M^s_{p,q}$ in the case $1 \leqslant p, q \leqslant \infty$, $s > 1/q'$ and he obtained that $\mathbb{Tr} M^s_{p,q}(\mathbb{R}^n) = M^{s-1/q'}_{p,q}(\mathbb{R}^{n-1})$. Frazier and Jawerth [8] proved that $\mathbb{Tr} B^s_{p,q}(\mathbb{R}^n) = B^{s-1/p}_{p,q}(\mathbb{R}^{n-1})$ in the case $0 < p, q \leqslant \infty$ and $s - 1/p > \max((n-1)(1/p - 1), 0)$.
In this paper, we will show the following:

**Theorem 1.2** *Let $n \geqslant 2$, $0 < p, q \leqslant \infty$, $s \in \mathbb{R}$. Then*

$$\mathbb{Tr} : f(x) \longrightarrow f(\bar{x}, 0), \; \bar{x} = (x_1, \cdots, x_{n-1}) \tag{1.3}$$

*is a retraction from $M^s_{p,q,\, p\wedge q\wedge 1}(\mathbb{R}^n)$ onto $M^s_{p,q}(\mathbb{R}^{n-1})$.*

Theorem 1.2 is sharp in the sense that $\mathbb{Tr} : M^s_{p,q,\,r}(\mathbb{R}^n) \not\to M^s_{p,q}(\mathbb{R}^{n-1})$ for some $r > 1$, $p, q \geqslant 1$. In view of the basic embedding $M^s_{p,q} \subset M^{s_2}_{p,q,q_2}$ for $s - s_2 > 1/q - 1/q_2 > 0$, $s \geqslant 0$, we immediately have

**Corollary 1.3** *Let $n \geqslant 2$, $0 < p, q \leqslant \infty$, $s \geqslant 0$. Let $\mathbb{T}\mathrm{r}$ be as in* (1.3). *Then for any $\varepsilon > 0$,*

$$\mathbb{T}\mathrm{r} : M_{p,q}^{s+\frac{1}{p\wedge q\wedge 1}-\frac{1}{q}+\varepsilon}(\mathbb{R}^n) \to M_{p,q}^{s}(\mathbb{R}^{n-1}).$$

One may ask if Corollary 1.3 holds for the limit case $s = \varepsilon = 0$, we can give a counterexample to show that $\mathbb{T}\mathrm{r} : M_{p,q}^{1/q'}(\mathbb{R}^n) \not\to M_{p,q}^{0}(\mathbb{R}^{n-1})$ in the case $p$, $q > 1$. Write

$$s_p = (n-1)(1/(p \wedge 1) - 1).$$

It is easy to see that $s_p = 0$ for $p \geqslant 1$ and $s_p = (n-1)(1/p - 1)$ for $p < 1$. For the trace of $\alpha$-modulation spaces, we have the following result.

**Theorem 1.4** *Let $n \geqslant 2$, $0 < p, q \leqslant \infty$, $s \geqslant \alpha(n-1)/q + \alpha s_p$. Let $\mathbb{T}\mathrm{r}$ be as in* (1.3). *Then*

$$\mathbb{T}\mathrm{r} : M_{p,p\wedge q\wedge 1}^{s+\alpha/p}(\mathbb{R}^n) \to M_{p,q}^{s}(\mathbb{R}^{n-1}).$$

The case $s < \alpha(n-1)/q + \alpha s_p$ is more complicated. We have the following

**Remark 1.5** *Let $n \geqslant 2$, $0 < p, q \leqslant \infty$, $s < \alpha(n-1)/q + \alpha s_p$. Let $\mathbb{T}\mathrm{r}$ be as in* (1.3). *Then*

$$\mathbb{T}\mathrm{r} : M_{p,p\wedge q\wedge 1}^{s+\sigma_{\alpha,p,q}}(\mathbb{R}^n) \to M_{p,q}^{s}(\mathbb{R}^{n-1}),$$

*where*

$$\sigma_{\alpha,p,q} = \begin{cases} \alpha/p + (1-\alpha)[\alpha(n-1)/q + \alpha s_p - s], & qs + (n-1)(1-\alpha) - q\alpha s_p > 0, \\ \alpha/p + \alpha s_p - s + \varepsilon, & qs + (n-1)(1-\alpha) - q\alpha s_p = 0, \\ \alpha/p + \alpha s_p - s, & qs + (n-1)(1-\alpha) - q\alpha s_p < 0. \end{cases}$$

Theorem 1.4 is sharp in the case $s \geqslant 0$, $p = q = 1$. As the end of this paper, we consider the trace of Besov spaces. If $s > s_p$, the corresponding result has been obtained in [8]. If $s \leqslant s_p$, we have the following trace theorem for Besov spaces:

**Theorem 1.6** *Let $n \geqslant 2$, $0 < p, q \leqslant \infty$, $s \leqslant s_p$. Let $\mathbb{T}\mathrm{r}$ be as in* (1.3). *Then we have*

$$\mathbb{T}\mathrm{r} : \tilde{B}_{p,p\wedge q\wedge 1}^{s_p+1/p}(\mathbb{R}^n) \to B_{p,q}^{s_p}(\mathbb{R}^{n-1}),$$

*and*

$$\mathbb{T}\mathrm{r} : B_{p,p\wedge q\wedge 1}^{s_p+1/p}(\mathbb{R}^n) \to B_{p,q}^{s}(\mathbb{R}^{n-1})$$

*in the case $s < s_p$. Moreover, when $1 < p < \infty$, we have*

$$\begin{aligned} &\mathbb{T}\mathrm{r} : \tilde{B}_{p,q\wedge 1}^{1/p,1/p}(\mathbb{R}^n) \to B_{p,q}^{0}(\mathbb{R}^{n-1}), \\ &\mathbb{T}\mathrm{r} : B_{p,q\wedge 1}^{1/p,s+1/p}(\mathbb{R}^n) \to B_{p,q}^{s}(\mathbb{R}^{n-1}), \qquad s < 0. \end{aligned}$$

The following are some notations which will be frequently used in this paper: $\mathbb{R}, \mathbb{N}$ and $\mathbb{Z}$ will stand for the sets of reals, positive integers and integers, respectively. $\mathbb{R}_+ = [0, \infty)$, $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. $c < 1$, $C > 1$ will denote positive universal constants, which can be different at different places. $a \lesssim b$ stands for $a \leqslant Cb$ for some constant $C > 1$, $a \sim b$ means that $a \lesssim b$ and $b \lesssim a$. We write $a \wedge b = \min(a, b)$, $a \vee b = \max(a, b)$. We denote by $p'$ the dual number of $p \in [1, \infty]$, i.e., $1/p + 1/p' = 1$. We will use Lebesgue spaces $L^p := L^p(\mathbb{R}^n)$, $\| \cdot \|_p := \| \cdot \|_{L^p}$, We denote by $\mathscr{S} := \mathscr{S}(\mathbb{R}^n)$ and $\mathscr{S}' := \mathscr{S}'(\mathbb{R}^n)$ the Schwartz space and tempered distribution space, respectively. $B(x, R)$ stands for the ball in $\mathbb{R}^n$ with center $x$ and radius $R$, $Q(x, R)$ denote the cube in $\mathbb{R}^n$ with center $x$ and side-length $2R$. $\mathscr{F}$ or $\widehat{\ }$ denotes the Fourier transform; $\mathscr{F}^{-1}$ denotes the inverse Fourier transform. For any set $A$ with finite elements, we denote by $\#A$ the number of the elements of $A$.

# 2 $\alpha$-modulation spaces

## 2.1 Definition

A countable set $\mathcal{Q}$ of subsets $Q \subset \mathbb{R}^n$ is said to be an admissible covering if $\mathbb{R}^n = \bigcup_{Q \in \mathcal{Q}} Q$ and there exists $n_0 < \infty$ such that $\#\{Q' \in \mathcal{Q} : Q \cap Q' \neq \varnothing\} \leqslant n_0$ for all $Q \in \mathcal{Q}$. Denote

$$\begin{aligned} r_Q &= \sup\{r \in \mathbb{R} : B(c_r, r) \subset Q\}, \\ R_Q &= \inf\{R \in \mathbb{R} : Q \subset B(c_R, R)\}. \end{aligned} \tag{2.1}$$

Let $0 \leqslant \alpha \leqslant 1$. An admissible covering is called an $\alpha$-covering of $\mathbb{R}^n$, if $|Q| \sim \langle x \rangle^{\alpha n}$ (uniformly) holds for all $Q \in \mathcal{Q}$ and for all $x \in Q$, and $\sup_{Q \in \mathcal{Q}} R_Q / r_Q \leqslant K$ for some $K < \infty$.

Let $\mathcal{Q}$ be an $\alpha$-covering of $\mathbb{R}^n$. A corresponding bounded admissible partition of unity of order $p$ ($p$-BAPU) $\{\psi_Q\}_{Q \in \mathcal{Q}}$ is a family of smooth functions satisfying

$$\begin{cases} \psi_Q : \mathbb{R}^n \to [0, 1], \quad \mathrm{supp} \psi_Q \subset Q, \\ \sum_{Q \in \mathcal{Q}} \psi_Q(\xi) \equiv 1 \quad \forall \xi \in \mathbb{R}^n, \\ \sup_{Q \in \mathcal{Q}} |Q|^{1/(p \wedge 1) - 1} \| \mathscr{F}^{-1} \psi_Q \|_{L^{(p \wedge 1)}} < \infty. \end{cases}$$

**Definition 2.1** *Let $0 < p, q \leqslant \infty$, $s \in \mathbb{R}$, $0 \leqslant \alpha \leqslant 1$. Let $\mathcal{Q}$ be an $\alpha$-covering of $\mathbb{R}^n$ with the p-BAPU $\{\psi_Q\}_{Q \in \mathcal{Q}}$. We denote by $M^{s,\alpha}_{p,q}$ the space of all tempered distributions $f$ for which the following is finite:*

$$\|f\|_{M^{s,\alpha}_{p,q}} = \left( \sum_{Q \in \mathcal{Q}} \langle \xi_Q \rangle^{qs} \| \mathscr{F}^{-1} \psi_Q \mathscr{F} f \|^q_{L^p(\mathbb{R}^n)} \right)^{1/q},$$

*where $\xi_Q \in Q$ is arbitrary. For $q = \infty$, we have a usual substitution for the $\ell^q$ norm with the $\ell^\infty$ norm.*

We now give an exact equivalent norm on $M^{s,\alpha}_{p,q}$. Denote

$$Q_k = Q(|k|^{\frac{\alpha}{1-\alpha}} k,\, r\langle k\rangle^{\frac{\alpha}{1-\alpha}}), \quad k \in \mathbb{Z}^n.$$

It is known that, there exists a constant $r_1 > 0$ such that for any $r > r_1$, $\{Q_k\}_{k\in\mathbb{Z}^n}$ is an $\alpha$-covering of $\mathbb{R}^n$, i.e., $\mathbb{R}^n = \bigcup_{k\in\mathbb{Z}^n} Q_k$ and there exists $n_0 \in \mathbb{N}$ such that $\#\{l \in \mathbb{Z}^n : Q_k \cap Q_{k+l} \neq \emptyset\} \leqslant n_0$. Moreover, $|Q_k| \sim \langle k\rangle^{\frac{n\alpha}{1-\alpha}}$. Let $\eta : \mathbb{R} \to [0,1]$ be a smooth bump function satisfying

$$\eta(\xi) := \begin{cases} 1, & |\xi| \leqslant 1, \\ \text{smooth}, & 1 < |\xi| \leqslant 2, \\ 0, & |\xi| \geqslant 2. \end{cases} \tag{2.2}$$

We write for $k = (k_1, ..., k_n)$ and $\xi = (\xi_1, ..., \xi_n)$,

$$\phi_{k_i}(\xi_i) = \eta\left(\frac{\xi_i - |k|^{\frac{\alpha}{1-\alpha}} k_i}{r\langle k\rangle^{\frac{\alpha}{1-\alpha}}}\right).$$

Put

$$\psi_k(\xi) = \frac{\phi_{k_1}(\xi_1)...\phi_{k_n}(\xi_n)}{\sum_{k\in\mathbb{Z}^n} \phi_{k_1}(\xi_1)...\phi_{k_n}(\xi_n)}, \quad k \in \mathbb{Z}^n. \tag{2.3}$$

We have

**Lemma 2.2** *Let $0 \leqslant \alpha < 1$, $0 < p \leqslant \infty$ and $\{\psi_k\}_{k\in\mathbb{Z}^n}$ be as in (2.3). Then $\{\psi_k\}_{k\in\mathbb{Z}^n}$ is a p-BAPU for $r > r_1$. In the case $\alpha = 0$, we can take $r_1 = 1/2$.*

**Proposition 2.3** *Let $0 \leqslant \alpha < 1$, $0 < p, q \leqslant \infty$, then*

$$\|f\|_{M^{s,\alpha}_{p,q}} = \left(\sum_{k\in\mathbb{Z}^n} \langle k\rangle^{\frac{qs}{1-\alpha}} \|\mathscr{F}^{-1}\psi_k \mathscr{F} f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}$$

*is an equivalent norm on $\alpha$-modulation space with the usual modification for $q = \infty$.*

**Proof.** See [1].

## 2.2 Equivalent norm via a New p-BAPU

We now construct a new covering, which is of importance for the proof of Theorem 1.4. Let $j \in \mathbb{Z}\backslash\{0\}$. We divide $[-|j|^{\frac{1}{1-\alpha}}, |j|^{\frac{1}{1-\alpha}}]$ into $2|j|/\{r_1\} = 2N_j$ intervals with equal length:

$$[-|j|^{\frac{1}{1-\alpha}},\, |j|^{\frac{1}{1-\alpha}}] = [r_{j,-N_j}, r_{j,-N_j+1}] \cup ... \cup [r_{j,N_j-1}, r_{j,N_j}].$$

Denote

$$\mathscr{R} = \{r_{j,s} : j \in \mathbb{N}, s = -N_j, \cdots, N_j\}.$$

We further write

$$\mathscr{K}_j^n = \{k = (k_1, \cdots, k_n) : k_i \in \mathscr{R},\, \max_{1\leqslant i\leqslant n} |k_i| = |j|^{\frac{1}{1-\alpha}}\}.$$

For any $k \in \mathscr{K}_j^n$, we write

$$Q_{kj} = Q(k, r|j|^{\frac{\alpha}{1-\alpha}}), \quad Q_{k0} = Q(0, 2).$$

We will write $\mathscr{K}_j = \mathscr{K}_j^n$ if there is no confusion.

**Proposition 2.4** *There exists $r_1 > 0$ such that for any $r > r_1$, $\{Q_{kj}\}_{k\in\mathscr{K}_j, j\in\mathbb{Z}_+}$ is an $\alpha$-covering of $\mathbb{R}^n$.*

**Proof.** Let $j \in \mathbb{N} \cup \{0\}$. We see that there exists $r_1 > 0$ such that for any $r > r_1$, $\{Q(|j|^{\frac{\alpha}{1-\alpha}} j, r|j|^{\frac{\alpha}{1-\alpha}})\}_j$ is an $\alpha$-covering of $\mathbb{R}$. Hence we easily see that

$$\mathbb{R} \subset \cup_{k\in\mathscr{K}_j, j\in\mathbb{Z}_+} Q_{kj}, \quad |Q_{kj}| \sim |j|^{\frac{n\alpha}{1-\alpha}} \sim \langle \xi_{Q_{kj}} \rangle^{n\alpha}, \forall\, \xi_{Q_{kj}} \in Q_{k_j},$$

$$\#\{Q_{k'j'} : Q_{kj} \cap Q_{k'j'} \neq \emptyset\} \leqslant n_0 < \infty.$$

Now, on the basis of the $\alpha$-covering constructed above, we further construct a $p$-BAPU. Let $j$ be fixed. Denote for $i = 1, \cdots, n$,

$$\phi_{kj}(\xi_i) = \phi\left(\frac{\xi_i - k_i}{r\langle j\rangle^{\frac{\alpha}{1-\alpha}}}\right), \quad k = (k_1, \cdots, k_n) \in \mathscr{K}_j.$$

$$\phi_{kj}(\xi) = \phi_{kj}(\xi_1)...\phi_{kj}(\xi_n).$$

We put

$$\psi_{kj}(\xi) = \frac{\phi_{kj}(\xi)}{\sum_{k\in\mathscr{K}_j, j\in\mathbb{Z}_+} \phi_{kj}(\xi)}. \tag{2.4}$$



Figure 1: $\alpha$-covering, the case of $n = 2$, $\alpha = 1/2$, $r_1 = 1$.

**Proposition 2.5** *Let* $0 < p < \infty$, $\psi_{kj}$ *be as in* (2.4). *Then* $\{\psi_{kj}\}_{k\in\mathscr{K}_j, j\in\mathbb{Z}_+}$ *is a* $p$*-BAPU.*

Noticing that $|\xi| \sim |j|^{1/(1-\alpha)}$ if $\xi \in \mathscr{K}_j$, $j \neq 0$, we immediately have

**Proposition 2.6** *Let* $0 < \alpha < 1$, $0 < p, q \leqslant \infty$, *then*

$$\|f\|_{M^{s,\alpha}_{p,q}} = \left( \sum_{j\in\mathbb{Z}_+} \langle j \rangle^{sq/(1-\alpha)} \sum_{k\in\mathscr{K}_j} \|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q}$$

*is another equivalent norm on* $\alpha$*-modulation space.*

## 2.3 Modulation spaces

In the case $\alpha = 0$, we get an equivalent norm on modulation spaces $M^s_{p,q}$:

$$\|f\|_{M^s_{p,q}} = \left( \sum_{k\in\mathbb{Z}^n} \langle k \rangle^{qs} \|\mathscr{F}^{-1}\psi_k\mathscr{F}f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q}. \tag{2.5}$$

The modulation spaces $M^s_{p,q}$ in the case $0 < p, q < 1$ was studied in [17, 18, 19] by using the norm (2.5). Soon after, Kobayashi [11] independently considered such a generalization in the case $0 < p, q < 1$.

Recalling that $\bar{x} = (x_1, ..., x_{n-1})$, we also define the following anisotropic modulation spaces $M^s_{p,q,r}$ for which the norm is defined as

$$\|f\|_{M^s_{p,q,r}} = \left( \sum_{k_n \in \mathbb{Z}} \left( \sum_{\bar{k} \in \mathbb{Z}^{n-1}} \langle \bar{k} \rangle^{qs} \|\mathscr{F}^{-1} \psi_k \mathscr{F} f\|^q_{L^p(\mathbb{R}^n)} \right)^{r/q} \right)^{1/r}.$$

This anisotropic version is of importance for the trace of modulation spaces.

## 2.4 Besov spaces

Write $\varphi(\cdot) = \eta(\cdot) - \eta(2\cdot)$ and $\varphi_k := \varphi(2^{-k}\cdot)$ for $k \geqslant 1$. $\varphi_0 := 1 - \sum_{k \geqslant 1} \varphi_k$. For simplicity, we write $\Delta_k = \mathscr{F}^{-1} \varphi_k \mathscr{F}$. The norm on Besov spaces $B^s_{p,q}(\mathbb{R}^n)$ are defined as follow:

$$\|f\|_{B^s_{p,q}} = \left( \sum_{j=0}^{\infty} 2^{sjq} \|\Delta_j f\|^q_p \right)^{1/q}.$$

For our purpose, we also need the following

$$\tilde{B}^s_{p,q}(\mathbb{R}^n) = \left( \sum_{k=0}^{\infty} k 2^{skq} \|\Delta_k f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q},$$

In the case $1 < p < \infty$, using Lizorkin's decomposition of $\mathbb{R}^n$, we have an equivalent quasi-norm on $B^s_{p,q}(\mathbb{R}^n)$. Let

$$K_k = \{x : |x_j| < 2^k, j = 1, 2, \dots, n\} \setminus \{x : |x_j| < 2^{k-1}, j = 1, 2, \dots, n\}$$

where $k \in \mathbb{Z}^+$ and

$$K_0 = \{x : |x_j| \leqslant 1, j = 1, 2, \dots, n\}$$

Subdivide $K_k$ with $k = 1, 2, 3 \dots$, by the $3n$ hyper-planes $\{x : x_m = 0\}$ and $\{x : x_m = \pm 2^{k-1}\}$, where $m = 1, \dots, n$, into cubes $P_{k,t}$. If $k$ is fixed, we obtain $T = 4^n - 2^n$ cubes. The cubes near the $n$-th axis are numbered by $t = 1, \dots, 2^n$ in an arbitrary way and the others are numbered by $t = 2^n + 1, \dots, T$. Let $P_{0,t} = K_0$, if $t = 1, \dots, T$. Then

$$\mathbb{R}_n = \cup_{k=0}^{\infty} \bar{K}_k = \cup_{k=0}^{\infty} \cup_{t=1}^{T} \bar{P}_{k,t}.$$



Figure 2: 1-covering, the case $n = 2$.

Let $\chi_{k,t}$ be a characteristic function on $P_{k,t}$. Then

$$\|f\|_{B^s_{p,q}(\mathbb{R}^n)} \asymp \left( \sum_{k=0}^{\infty} \sum_{t=1}^{T} 2^{skq} \|\mathscr{F}^{-1}\chi_{k,t}\mathscr{F} f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q}.$$

We construct two new norms. For simplicity, we write

$$\Delta_{k,t} = \mathscr{F}^{-1}\chi_{k,t}\mathscr{F}.$$

Define

$$B^{s_1,s_2}_{p,q} = \left( \sum_{k=0}^{\infty} \left( \sum_{t=1}^{2^n} 2^{s_1kq} \|\Delta_{k,t} f\|^q_{L^p(\mathbb{R}^n)} + \sum_{t=2^n+1}^{T} 2^{s_2kq} \|\Delta_{k,t} f\|^q_{L^p(\mathbb{R}^n)} \right) \right)^{1/q},$$

$$\tilde{B}^{s_1,s_2}_{p,q} = \left( \sum_{k=0}^{\infty} \left( \sum_{t=1}^{2^n} k 2^{s_1kq} \|\Delta_{k,t} f\|^q_{L^p(\mathbb{R}^n)} + \sum_{t=2^n+1}^{T} 2^{s_2kq} \|\Delta_{k,t} f\|^q_{L^p(\mathbb{R}^n)} \right) \right)^{1/q}.$$

# 3 Proof of Theorem 1.2

If there is no explanation, we always assume $r = 1/2$ in the $p$-BAPU for the case of modulation spaces. To show our main theorem, we will use the following

**Lemma 3.1** (Triebel, [14]) *Let* $\Omega$ *be a compact subset of* $\mathbb{R}^n$ *and* $0 < p \leqslant \infty$. *Denote* $L_p^\Omega = \{f \in L^p : Supp\hat{f} \subset \Omega\}$. *Let* $0 < r < p$. *Then*

$$\left\| \sup_{z \in \mathbb{R}^n} \frac{f(\cdot - z)}{1 + |z|^{n/r}} \right\|_{L^p(\mathbb{R}^n)} \lesssim \|f\|_{L^p(\mathbb{R}^n)},$$

*holds for any* $f \in L_p^\Omega$.

Assume that $\text{Supp}\hat{f} \subset B(\xi_0, R)$. It is easy to see that for $g = e^{ix\xi_0} f(R^{-1}\cdot)$, $\hat{g} = R^n \hat{f}(R(\xi - \xi_0))$. It follows that $\text{Supp}\hat{g} \subset B(0,1)$. Taking $\Omega = B(0,1)$ in Lemma 3.1, we find that

$$\left\| \sup_{z \in \mathbb{R}^n} \frac{g(\cdot - z)}{1 + |z|^{n/r}} \right\|_{L^p(\mathbb{R}^n)} \lesssim \|g\|_{L^p(\mathbb{R}^n)},$$

By scaling, we have

$$\left\| \sup_{z \in \mathbb{R}^n} \frac{f(\cdot - z)}{1 + |Rz|^{n/r}} \right\|_{L^p(\mathbb{R}^n)} \leqslant C\|f\|_{L^p(\mathbb{R}^n)} \tag{3.1}$$

Note that the constant $C$ in (3.1) is independent of $f \in L^p_{B(\xi_0,R)} = \{f \in L^p : \text{Supp}\hat{f} \subset B(\xi_0, R)\}$. It is also independent of $\xi_0 \in \mathbb{R}^n$.

For convenience, we write

$$\Box_k = \mathscr{F}^{-1}\psi_k\mathscr{F}, \quad k \in \mathbb{Z}^n.$$

We define the maximum function $M_k^* f$ as follows:

$$M_k^* f = \sup_{y \in \mathbb{Z}^n} \frac{|\Box_k f(x - y)|}{1 + |y|^{n/r}}. \tag{3.2}$$

Taking $y_1 = ... = y_{n-1} = 0$, $y_n = x_n$ in (3.2), we have for $|x_n| \leqslant 1$,

$$|(\Box_k f)(\bar{x}, 0)| \lesssim |M_k^* f(x)|, \quad \bar{x} = (x_1, \cdots, x_{n-1})$$

Hence

$$\|(\Box_k f)(\cdot, 0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim \|M_k^* f(\cdot, x_n)\|_{L^p(\mathbb{R}^{n-1})}, \tag{3.3}$$

Integrating (3.3) over $x_n \in [0,1]$, one has that

$$\|(\Box_k f)(\cdot, 0)\|^p_{L^p(\mathbb{R}^{n-1})} \lesssim \int_{\mathbb{R}} \|M_k^* f(\cdot, x_n)\|^p_{L^p(\mathbb{R}^{n-1})} dx_n,$$

Hence

$$\|(\Box_k f)(\cdot, 0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim \|M_k^* f\|_{L^p(\mathbb{R}^n)}. \tag{3.4}$$

We denote by $\mathscr{F}_{\bar{x}}$ ($\mathscr{F}^{-1}_{\bar{\xi}}$) the partial (inverse) Fourier transform on $\bar{x}$ ($\bar{\xi}$). Write $\psi_{\bar{k}}(\bar{x})$ as the $p$-BAPU functions in $\mathbb{R}^{n-1}$ as in (2.3), i.e.,

$$\psi_{\bar{k}}(\bar{\xi}) = \frac{\phi_{k_1}(\xi_1)...\phi_{k_{n-1}}(\xi_{n-1})}{\sum_{\bar{k}\in\mathbb{Z}^{(n-1)}} \phi_{k_1}(\xi_1)\cdots\phi_{k_{n-1}}(\xi_{n-1})}, \quad \bar{k} \in \mathbb{Z}^{(n-1)}. \tag{3.5}$$

Then we have

$$\begin{aligned}(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}\mathscr{F}_{\bar{x}})(\bar{x},0) &= \sum_{l\in\mathbb{Z}^n}(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}\mathscr{F}_{\bar{x}}\mathscr{F}^{-1}\psi_l\mathscr{F})(\bar{x},0)\\ &= \sum_{l\in\mathbb{Z}^n}(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}) * (\mathscr{F}^{-1}\psi_l\mathscr{F})(\bar{x},0)\end{aligned}$$

From the support property of $\psi_l$ as in (2.3), we find that

$$\psi_{\bar{k}}\psi_l = 0, \quad \text{if } |\bar{l}-\bar{k}| \geqslant C.$$

Hence

$$(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}\mathscr{F}_{\bar{x}})(\bar{x},0) = \sum_{l\in\mathbb{Z}^n,\ |\bar{k}-\bar{l}|\leqslant C}(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}) * ((\mathscr{F}^{-1}\psi_l\mathscr{F})(\cdot,0))$$

*Case 1.* $1 \leqslant p \leqslant \infty$. Using Young's inequality, (3.1) and (3.4), we obtain

$$\begin{aligned}&\|\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}\mathscr{F}_{\bar{x}}f(\bar{x},0)\|_{L^p(\mathbb{R}^{n-1})}\\ &\lesssim \sum_{l\in\mathbb{Z}^n,|\bar{k}-\bar{l}|\leqslant C}\|\mathscr{F}^{-1}_{\bar{\xi}}\psi_{\bar{k}}\|_{L^1(\mathbb{R}^{n-1})}\|\mathscr{F}^{-1}\psi_l\mathscr{F}f\|_{L^p(\mathbb{R}^{n-1})}\\ &\lesssim \sum_{l\in\mathbb{Z}^n,|\bar{k}-\bar{l}|\leqslant C}\|M^*_l f\|_{L^p(\mathbb{R}^n)}\\ &\lesssim \sum_{l\in\mathbb{Z}^n,|\bar{k}-\bar{l}|\leqslant C}\|\Box_l f\|_{L^p(\mathbb{R}^n)}.\end{aligned}$$

Hence,

$$\|f(\bar{x},0)\|_{M^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left(\sum_{\bar{k}\in\mathbb{Z}^{n-1}}\langle\bar{k}\rangle^{sq}\left(\sum_{l\in\mathbb{Z}^n,|\bar{k}-\bar{l}|\leqslant C}\|\Box_l f\|_{L^p(\mathbb{R}^n)}\right)^q\right)^{1/q}.$$

If $0 < q \leqslant 1$, then

$$\begin{aligned}\|f(\bar{x},0)\|_{M^s_{p,q}(\mathbb{R}^{n-1})} &\lesssim \left(\sum_{l\in\mathbb{Z}^n}\sum_{\bar{k}\in\mathbb{Z}^{n-1}}\langle\bar{k}\rangle^{sq}\chi_{(|\bar{k}-\bar{l}|\leqslant C)}\|\Box_l f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}\\ &\lesssim \left(\sum_{l\in\mathbb{Z}^n}\langle\bar{l}\rangle^{sq}\|\Box_l f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q} = \|f\|_{M^s_{p,q,q}}.\end{aligned}$$

If $1 \leqslant q \leqslant \infty$, using Minkowski's inequality together with Hölder's inequality,

$$\begin{aligned}
\|f(\cdot,0)\|_{M^s_{p,q}(\mathbb{R}^{n-1})} &\lesssim \left( \sum_{\bar{k},\bar{l}\in\mathbb{Z}^{n-1}} \langle \bar{k}\rangle^{sq} \left( \sum_{l_n\in\mathbb{Z}} \chi_{(|\bar{k}-\bar{l}|\leqslant C)} \|\Box_l f\|_{L^p(\mathbb{R}^n)} \right)^q \right)^{1/q} \\
&\lesssim \sum_{l_n\in\mathbb{Z}} \left( \sum_{\bar{k},\bar{l}\in\mathbb{Z}^{n-1}} \langle \bar{k}\rangle^{sq} \chi_{(|\bar{k}-\bar{l}|\leqslant C)} \|\Box_l f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q} \\
&\lesssim \sum_{l_n\in\mathbb{Z}} \left( \sum_{\bar{l}\in\mathbb{Z}^{n-1}} \langle \bar{l}\rangle^{sq} \|\Box_l f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q} \\
&= \|f\|_{M^s_{p,q,1}}.
\end{aligned}$$

To begin with the proof for the case $0 < p < 1$, we need the following lemma:

**Lemma 3.2** *Let* $0 < p \leqslant 1$. *Suppose that* $f, g \in L^p_{B(x_0,R)}$, *then there exists a constant* $C > 0$ *which is independent of* $x_0 \in \mathbb{R}^n$ *and* $R > 0$ *such that*

$$\|f * g\|_p \leqslant C R^{n(\frac{1}{p}-1)} \|f\|_p \|g\|_p.$$

**Proof.** In the case $f, g \in L^p_{B(0,1)}$, we have

$$\|f * g\|_p \lesssim \|f\|_p \|g\|_p.$$

Taking $f_\lambda = f(\lambda\cdot)$ and $g_\lambda = g(\lambda\cdot)$, we see that

$$f_{R^{-1}}, g_{R^{-1}} \in L^p_{B(0,1)}, \quad \text{if } f, g \in L^p_{B(0,R)}.$$

Hence, for any $f, g \in L^p_{B(0,R)}$,

$$\|f_{R^{-1}} * g_{R^{-1}}\|_p \lesssim \|f_{R^{-1}}\|_p \|g_{R^{-1}}\|_p.$$

By scaling, we have

$$\|f * g\|_p \lesssim R^{n(\frac{1}{p}-1)} \|f\|_p \|g\|_p.$$

By a translation $\widehat{e^{ix_0x}f} = \hat{f}(\xi - x_0)$, we immediately have the result, as desired. □

*Case 2.* $0 < p < 1$. By Lemma 3.2, (3.1) and (3.4),

$$\begin{aligned}
&\|\mathscr{F}^{-1}_{\bar{\xi}} \psi_{\bar{k}} \mathscr{F}_{\bar{x}} f(\bar{x},0)\|^p_{L^p(\mathbb{R}^{n-1})} \\
&\lesssim \sum_{l\in\mathbb{Z}^n, |\bar{k}-\bar{l}|\leqslant C} \|\mathscr{F}^{-1}_{\bar{\xi}} \psi_{\bar{k}}\|^p_{L^p(\mathbb{R}^{n-1})} \|(\mathscr{F}^{-1}\psi_l \mathscr{F} f)(\cdot,0)\|^p_{L^p(\mathbb{R}^{n-1})}
\end{aligned}$$

$$
\begin{aligned}
&\lesssim \sum_{l\in\mathbb{Z}^n,|\bar{k}-\bar{l}|\leqslant C} \|(\mathscr{F}^{-1}\psi_l\mathscr{F}f)(\cdot,0)\|^p_{L^p(\mathbb{R}^{n-1})}\\
&\lesssim \sum_{l\in\mathbb{Z}^n,|\bar{k}-\bar{l}|\leqslant C} \|\Box_l f\|^p_{L^p(\mathbb{R}^n)}.
\end{aligned}
$$

It follows that

$$
\|f(\bar{x},0)\|_{M^{s,\alpha}_{p,q}} \lesssim \left(\sum_{\bar{k}\in\mathbb{Z}^n}\langle\bar{k}\rangle^{sq}\left(\sum_{l\in\mathbb{Z}^n}\|\Box_l f\|^p_{L^p(\mathbb{R}^n)}\chi_{(|\bar{k}-\bar{l}|\leqslant C)}\right)^{q/p}\right)^{1/q}.
$$

If $q\leqslant p$, one has that

$$
\begin{aligned}
\|f(\bar{x},0)\|_{M^s_{p,q}(\mathbb{R}^{n-1})} &\lesssim \left(\sum_{l\in\mathbb{Z}^n}\sum_{k\in\mathbb{Z}^n}\langle\bar{k}\rangle^{sq}\|\Box^\alpha_l f\|^q_{L^p(\mathbb{R}^n)}\chi_{(|\bar{k}-\bar{l}|\leqslant C)}\right)^{1/q}\\
&\lesssim \|f\|_{M^s_{p,q}}.
\end{aligned}
$$

If $q\geqslant p$, using Minkowski's inequality, we have

$$
\begin{aligned}
\|f(\bar{x},0)\|_{M^s_{p,q}(\mathbb{R}^{n-1})} &\lesssim \left(\sum_{l_n\in\mathbb{Z}}\left(\sum_{\bar{k},\bar{l}\in\mathbb{Z}^{n-1}}\langle\bar{k}\rangle^{sq}\|\Box_l f\|^q_{L^p(\mathbb{R}^n)}\chi_{(|\bar{k}-\bar{l}|\leqslant C)}\right)^{p/q}\right)^{1/p}\\
&\lesssim \left(\sum_{l_n\in\mathbb{Z}}\left(\sum_{\bar{l}\in\mathbb{Z}^{n-1}}\langle\bar{l}\rangle^{sq}\|\Box_l f\|^q_{L^p(\mathbb{R}^n)}\right)^{p/q}\right)^{1/p}\\
&= \|f\|_{M^s_{p,q,p}}.
\end{aligned}
$$

In order to show $\mathbb{Tr}$ is a retraction, we need to show the existence of $\mathbb{Tr}^{-1}$. Let $\eta$ be as in (2.2) satisfying $(\mathscr{F}^{-1}_{\xi_n}\eta)(0)=1$. For any $f\in M^s_{p,q}(\mathbb{R}^{n-1})$, we define

$$
g(x)=[(\mathscr{F}^{-1}_{\xi_n}\eta)(x_n)]f(\bar{x}):=(\mathbb{Tr}^{-1}f)(x).
$$

It is easy to see that $g(\bar{x},0)=f(\bar{x})$ and $\Box_k g=0$ for $|k_n|\geqslant 3$. Hence,

$$
\begin{aligned}
\|g\|_{M^s_{p,q,p\wedge q\wedge 1}(\mathbb{R}^n)} &\lesssim \left(\sum_{k_n\in\mathbb{Z}}\left(\sum_{\bar{k}\in\mathbb{Z}^{n-1}}\langle\bar{k}\rangle^{sq}\|\Box_k g\|^q_{L^p(\mathbb{R}^n)}\right)^{p\wedge q\wedge 1/q}\right)^{1/p\wedge q\wedge 1}\\
&= \sum_{|k_n|\leqslant 2}\left(\sum_{\bar{k}\in\mathbb{Z}^{n-1}}\langle\bar{k}\rangle^{sq}\|\Box_{\bar{k}}f\|^q_{L^p(\mathbb{R}^n)}\|\mathscr{F}^{-1}_{\xi_n}\eta\|^q_{L^p(\mathbb{R})}\right)^{1/q}
\end{aligned}
$$

$$\lesssim \|f\|_{M^s_{p,q}(\mathbb{R}^{n-1})}.$$

It follows that $\mathbb{Tr}^{-1}: M^s_{p,q}(\mathbb{R}^{n-1}) \to M^s_{p,q,p\wedge q\wedge 1}(\mathbb{R}^n)$. $\square$

As the end of this section, we show that Theorem 1.2 and Corollary 1.3 are sharp conclusions. First, we show that $\mathbb{Tr}: M^0_{p,q,r}(\mathbb{R}^n) \not\to M^0_{p,q}(\mathbb{R}^{n-1})$ if $r > 1$. Let $\eta$ be as in (2.2), $f = \mathscr{F}^{-1}(\eta(2\xi_1)...\eta(2\xi_n))$. For $k = (k_1, ..., k_n)$, we denote

$$F(x) = \sum_{|k_n|\leqslant 2^N} \langle k_n\rangle^{-1} e^{\mathrm{i}k_n x_n} f(x).$$

It is easy to see that

$$\mathscr{F}F(\xi) = \sum_{|k_n|\leqslant 2^N} \langle k_n\rangle^{-1} \eta(2\xi_1)...\eta(2(\xi_n - k_n)).$$

Hence, $\square_k F = 0$ if $\max_{i=1,...,n-1}|k_i| > 2$ or $|k_n| > 2^N+1$. In view of the definition

$$\begin{aligned}
\|F\|_{M^0_{p,q,r}(\mathbb{R}^n)} &\lesssim \left(\sum_{|k_n|\leqslant 2^N+1}\left(\sum_{|k_i|\leqslant 2,\, 1\leqslant i\leqslant n-1} \|\square_k F\|^q_{L^p(\mathbb{R}^n)}\right)^{r/q}\right)^{1/r} \\
&\lesssim \sum_{|k_i|\leqslant 2,\, 1\leqslant i\leqslant n-1}\left(\sum_{|k_n|\leqslant 2^N+1} \|\square_k F\|^r_{L^p(\mathbb{R}^n)}\right)^{1/r} \\
&\lesssim \left(\sum_{|k_n|\leqslant 2^N+1} \langle k_n\rangle^{-r}\right)^{1/r} \lesssim 1.
\end{aligned}$$

On the other hand, we may assume that $(\mathscr{F}^{-1}_{\xi_n}\eta(2\cdot))(0) = 1$. We have

$$F(\bar{x}, 0) = \left(\sum_{|k_n|\leqslant 2^N} \langle k_n\rangle^{-1}\right) \mathscr{F}^{-1}_{\bar{\xi}}[\eta(2\xi_1)...\eta(2(\xi_{n-1}))].$$

So,

$$\begin{aligned}
\|F\|_{M^0_{p,q}(\mathbb{R}^{n-1})} &\gtrsim \left(\sum_{|k_n|\leqslant 2^N} \langle k_n\rangle^{-1}\right)\left(\sum_{|k_i|\leqslant 2,\, 1\leqslant i\leqslant n-1} \|\mathscr{F}^{-1}\psi_{\bar{k}}\eta(2\xi_1)...\eta(2\xi_{n-1})\|^q_{L^p(\mathbb{R}^{n-1})}\right)^{1/q} \\
&\gtrsim N.
\end{aligned}$$

Let $N \to \infty$, we have $\mathbb{Tr}: M^0_{p,q,r}(\mathbb{R}^n) \not\to M^0_{p,q}(\mathbb{R}^{n-1})$.

Next, we show that $\mathbb{T}\mathrm{r} : M_{p,q}^{1/q'}(\mathbb{R}^n) \not\to M_{p,q}^0(\mathbb{R}^{n-1})$ as $q > 1$. For $k = (k_1, ..., k_n)$, we denote

$$F(x) = \sum_{|k_n| \leqslant 2^N} \frac{1}{\langle k_n \rangle \ln \langle k_n \rangle} e^{\mathrm{i} k_n x_n} f(x).$$

Similarly as in the above, we have

$$\begin{aligned} \|F\|_{M_{p,q}^{1/q'}(\mathbb{R}^n)} &\lesssim \left( \sum_{|k_n| \leqslant 2^N+1} \sum_{|k_i| \leqslant 2,\, 1 \leqslant i \leqslant n-1} \langle k_n \rangle^{q-1} \|\Box_k F\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q} \\ &\lesssim \left( \sum_{|k_n| \leqslant 2^N+1} \frac{1}{\langle k_n \rangle \ln^q \langle k_n \rangle} \right)^{1/q} \lesssim 1. \end{aligned}$$

On the other hand,

$$\|F\|_{M_{p,q}^0(\mathbb{R}^{n-1})} \gtrsim \left( \sum_{|k_n| \leqslant 2^N} \frac{1}{\langle k_n \rangle \ln \langle k_n \rangle} \right) \to \infty, \quad N \to \infty.$$

# 4 Proof of Theorem 1.4 and Remark 1.5

For convenience, we write

$$\Box_{k,j}^\alpha = \mathscr{F}^{-1} \psi_{k,j} \mathscr{F}, \quad k \in \mathscr{K}_j,\ j \in \mathbb{Z}_+.$$

We define the maximum function $M_{k,j}^* f$ as follows:

$$M_{k,j}^* f = \sup_{y \in \mathbb{Z}^n} \frac{|\Box_{k,j}^\alpha f(x-y)|}{1 + |\langle j \rangle^{\alpha/(1-\alpha)} y|^{n/r}}. \tag{4.1}$$

Taking $y_1 = ... = y_{n-1} = 0$, $y_n = x_n$ in (4.1), we have for $\langle j \rangle^{-\alpha/(1-\alpha)} \leqslant |x_n| \leqslant 2\langle j \rangle^{-\alpha/(1-\alpha)}$,

$$|(\Box_{k,j}^\alpha f)(\bar{x}, 0)| \lesssim |M_{k,j}^* f(x)|, \quad \bar{x} = (x_1, \cdots, x_{n-1}).$$

Hence

$$\|(\Box_{k,j}^\alpha f)(\cdot, 0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim \|M_{k,j}^* f(\cdot, x_n)\|_{L^p(\mathbb{R}^{n-1})}, \tag{4.2}$$

Integrating (4.2) over $x_n \in [\langle j \rangle^{-\alpha/(1-\alpha)}, 2\langle j \rangle^{-\alpha/(1-\alpha)}]$, one has that

$$\|(\Box_{k,j}^\alpha f)(\cdot, 0)\|^p_{L^p(\mathbb{R}^{n-1})} \lesssim \langle j \rangle^{\alpha/(1-\alpha)} \int_{\mathbb{R}} \|M_{k,j}^* f(\cdot, x_n)\|^p_{L^p(\mathbb{R}^{n-1})} dx_n,$$

Hence

$$\|(\square^{\alpha}_{k,j}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim \langle j\rangle^{\alpha/p(1-\alpha)}\|M^*_{k,j}f\|_{L^p(\mathbb{R}^n)}. \tag{4.3}$$

We denote by $\mathscr{F}_{\bar{x}}(\mathscr{F}^{-1}_{\bar{\xi}})$ the partial (inverse) Fourier transform on $\bar{x}=(x_1,...,x_{n-1})$ $(\bar{\xi}=(\xi_1,\cdots,\xi_{n-1}))$. Write $\psi_{m,l}(\bar{x})$ as the $p$-BAPU functions in $\mathbb{R}^{n-1}$ as in (2.4). So, by the definition,

$$\|f\|_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})} = \left(\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}^{n-1}_l}\langle l\rangle^{\frac{qs}{1-\alpha}}\|\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l}(\bar{\xi})\mathscr{F}_{\bar{x}}f)(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}\right)^{1/q}. \tag{4.4}$$

In order to have no confusion, we always denote by $\psi_{m,l}$ the $p$-BAPU function in $\mathbb{R}^{n-1}$ and by $\psi_{k,j}$ the $p$-BAPU function in $\mathbb{R}^n$. From the support property of $\psi_{k,j}$, we find that

$$(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l}\mathscr{F}_{\bar{x}}f)(\bar{x},0) = \sum_{j\geqslant l-C,\,k\in\mathscr{K}^n_j}(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l}\mathscr{F}_{\bar{x}}\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\bar{x},0). \tag{4.5}$$

For our purpose we further decompose $\mathscr{K}^n_j$. Denote

$$\mathscr{K}^n_{j,\lambda} = \{k\in\mathscr{K}^n_j:\ \max_{1\leqslant i\leqslant n-1}|k_i| = \lambda\}, \quad \lambda = r_{j0}, r_{j1}, ..., r_{jN_j}, \quad r_{j0}=0.$$

We easily see that $\sum_{k\in\mathscr{K}^n_j} = \sum_{\lambda=0,r_{j1},...,r_{jN_j}}\sum_{\mathscr{K}^n_{j,\lambda}}$, $N_j\sim\langle j\rangle$. Now we divide our discussion into the following four cases.

*Case 1.* $1\leqslant p\leqslant\infty$ and $0<q\leqslant 1$. By (4.4) and (4.5),

$$\begin{aligned}\|f(\cdot,0)\|^q_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})} \lesssim \sum_{j\in\mathbb{Z}_+}\sum_{\lambda=0,r_{j1},...,r_{jN_j}}\sum_{k\in\mathscr{K}^n_{j,\lambda}}\sum_{l\leqslant j+C}\sum_{m\in\mathscr{K}^{n-1}_l}\langle l\rangle^{\frac{sq}{1-\alpha}}\\ \times\|(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l})*(\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}.\end{aligned} \tag{4.6}$$

In order to control (4.6) by $\|f\|_{M^{s+\alpha/p,\alpha}_{p,q}}$, we need to bound the sum $\sum_{l\leqslant j+C}\sum_{m\in\mathscr{K}^{n-1}_l}$. It is easy to see that for fixed $k,j$,

$$\#\{m\in\mathscr{K}^{n-1}_l:\ \mathrm{supp}\,\psi_{m,l}\cap\mathrm{supp}\,\psi_{k,j}(\cdot,0)\neq\varnothing\} \lesssim \min\left(\langle l\rangle^{n-2},\ \langle j\rangle^{\frac{\alpha(n-2)}{1-\alpha}}/\langle l\rangle^{\frac{\alpha(n-2)}{1-\alpha}}\right). \tag{4.7}$$

Moreover, $k\in\mathscr{K}^n_{j,r_{ja}}$ means that $\mathrm{supp}\,\psi_{m,l}\cap\mathrm{supp}\,\psi_{k,j}(\cdot,0)\neq\varnothing$ only if $a^{1-\alpha}\langle j\rangle^{\alpha}\lesssim l\lesssim(1+a)^{1-\alpha}\langle j\rangle^{\alpha}$. Hence, in view of Young's inequality, (4.3),

$$\Delta_{ja} := \sum_{k\in\mathscr{K}^n_{j,r_{ja}}}\sum_{m\in\mathscr{K}^{n-1}_l,\,l\leqslant j+C}\langle l\rangle^{\frac{sq}{1-\alpha}}\|(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l})*(\mathscr{F}^{-1}\psi_{k,j}\mathscr{F})f(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}$$

$$\begin{aligned}
&\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \sum_{a^{1-\alpha}\langle j\rangle^{\alpha}\lesssim l\lesssim (1+a)^{1-\alpha}\langle j\rangle^{\alpha}} \langle l\rangle^{\frac{sq}{1-\alpha}} \min\Big(\langle l\rangle^{n-2},\ \langle j\rangle^{\frac{\alpha(n-2)}{1-\alpha}}/\langle l\rangle^{\frac{\alpha(n-2)}{1-\alpha}}\Big)\\
&\qquad \times \|(\mathscr{F}^{-1}\psi_{k,j}\mathscr{F})f(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}.\\
&\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \sum_{a^{1-\alpha}\langle j\rangle^{\alpha}\lesssim l\lesssim (1+a)^{1-\alpha}\langle j\rangle^{\alpha}} \langle l\rangle^{\frac{sq}{1-\alpha}} \min\Big(\langle l\rangle^{n-2},\ \langle j\rangle^{\frac{\alpha(n-2)}{1-\alpha}}/\langle l\rangle^{\frac{\alpha(n-2)}{1-\alpha}}\Big)\\
&\qquad \times \langle j\rangle^{\frac{q\alpha}{p(1-\alpha)}}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}. \qquad (4.8)
\end{aligned}$$

We discuss the following four subcases.

*Case 1A.* $\alpha(n-1)\leqslant qs$. If $a=0$, one has that

$$\begin{aligned}
\Delta_{ja} &\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \sum_{0\leqslant l\lesssim\langle j\rangle^{\alpha}} \langle l\rangle^{\frac{sq}{1-\alpha}+(n-2)}\langle j\rangle^{\frac{q\alpha}{p(1-\alpha)}}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}\\
&\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{q}{1-\alpha}\left(\alpha s+\frac{\alpha}{p}+(1-\alpha)\frac{\alpha(n-1)}{q}\right)}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}\\
&\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{q}{1-\alpha}\left(s+\frac{\alpha}{p}\right)}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}. \qquad (4.9)
\end{aligned}$$

If $a\geqslant 1$, we have

$$\begin{aligned}
\Delta_{ja} &\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \sum_{a^{1-\alpha}\langle j\rangle^{\alpha}\lesssim l\lesssim (1+a)^{1-\alpha}\langle j\rangle^{\alpha}} \langle l\rangle^{\frac{sq}{1-\alpha}-\frac{\alpha(n-2)}{1-\alpha}}\langle j\rangle^{\frac{q\alpha}{p(1-\alpha)}+\frac{\alpha(n-2)}{1-\alpha}}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}\\
&\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} a^{qs-\alpha(n-1)}\langle j\rangle^{\frac{\alpha sq}{1-\alpha}+\frac{q\alpha}{p(1-\alpha)}+\alpha(n-1)}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}. \qquad (4.10)
\end{aligned}$$

It follows from $qs\geqslant\alpha(n-1)$ that $\Delta_{ja}$ takes the maximal value as $a=N_j\sim\langle j\rangle$. Hence,

$$\Delta_{ja}\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{q}{1-\alpha}\left(s+\frac{\alpha}{p}\right)}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}. \qquad (4.11)$$

Inserting the estimates of $\Delta_{ja}$ as in (4.9) and (4.11) into (4.6), we have

$$\|f(\cdot,0)\|^q_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})}\lesssim \sum_{j\in\mathbb{Z}_+}\sum_{k\in\mathscr{K}^n_j}\langle j\rangle^{\frac{q}{1-\alpha}(s+\frac{\alpha}{p})}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}=\|f\|^q_{M^{s+\alpha/p,\alpha}_{p,q}}.$$

*Case 1B.* $\alpha(n-1)>qs$ and $qs+(1-\alpha)(n-1)>0$. If $a\geqslant 1$, from (4.10) and $\alpha(n-1)>qs$ we have

$$\Delta_{ja}\lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{\alpha sq}{1-\alpha}+\frac{q\alpha}{p(1-\alpha)}+\alpha(n-1)}\|\Box^{\alpha}_{k,j}f\|^q_{L^p(\mathbb{R}^n)}. \qquad (4.12)$$

Since $qs + (1-\alpha)(n-1) > 0$, similar to (4.9), we see that (4.12) also holds for the case $a = 0$. It follows that

$$\|f(\cdot,0)\|_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})} \lesssim \|f\|_{M_{p,q}^{\alpha s+(1-\alpha)\frac{\alpha(n-1)}{q}+\alpha/p,\,\alpha}(\mathbb{R}^n)}.$$

*Case 1C.* $qs = -(n-1)(1-\alpha)$. Using the first estimate as in (4.9), we have for $a = 0$,

$$\Delta_{ja} \lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{q\alpha}{p(1-\alpha)}} \ln\langle j\rangle \|\Box^\alpha_{k,j} f\|^q_{L^p(\mathbb{R}^n)}. \tag{4.13}$$

For $a \geqslant 1$,

$$\Delta_{ja} \lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{\alpha sq}{1-\alpha}+\frac{q\alpha}{p(1-\alpha)}+\alpha(n-1)} \|\Box^\alpha_{k,j} f\|^q_{L^p(\mathbb{R}^n)}. \tag{4.14}$$

This implies that

$$\|f(\cdot,0)\|_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})} \lesssim \|f\|_{M^{\alpha/p+\varepsilon,\,\alpha}_{p,q}(\mathbb{R}^n)}.$$

*Case 1D.* $qs < -(n-1)(1-\alpha)$. It is easy to see that $\ln\langle j\rangle$ can be removed in (4.13). So, we have the result, as desired.

*Case 2.* $1 \leqslant p, q \leqslant \infty$. Using Minkowski's inequality, we have

$$\begin{aligned}
&\|f(\cdot,0)\|_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})}\\
&\lesssim \left(\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}^{n-1}_l} \langle l\rangle^{\frac{sq}{1-\alpha}} \left(\sum_{j\in\mathbb{Z}_+}\sum_{k\in\mathscr{K}^n_j} \|(\mathscr{F}^{-1}_{\bar\xi}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F} f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}\right)^q\right)^{1/q}\\
&\lesssim \sum_{j\in\mathbb{Z}_+}\sum_{a=0}^{N_j}\sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \left(\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}^{n-1}_l} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(\mathscr{F}^{-1}_{\bar\xi}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F} f)(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}\right)^{1/q}.
\end{aligned} \tag{4.15}$$

Using the same way as in Case 1, we can get that for any $k \in \mathscr{K}^n_{j,r_{ja}}$,

$$\begin{aligned}
&\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}^{n-1}_l} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(\mathscr{F}^{-1}_{\bar\xi}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F} f)(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}\\
&\lesssim \sum_{a^{1-\alpha}\langle j\rangle^\alpha \lesssim l \lesssim (1+a)^{1-\alpha}\langle j\rangle^\alpha} \langle l\rangle^{\frac{sq}{1-\alpha}} \min\left(\langle l\rangle^{n-2}, \frac{\langle j\rangle^{\frac{\alpha(n-2)}{1-\alpha}}}{\langle l\rangle^{\frac{\alpha(n-2)}{1-\alpha}}}\right) \langle j\rangle^{\frac{q\alpha}{p(1-\alpha)}} \|\Box^\alpha_{k,j} f\|^q_{L^p(\mathbb{R}^n)}.
\end{aligned} \tag{4.16}$$

Repeating the calculation procedure as in Case 1, we have the result, as desired.

*Case 3.* $0 < q \leqslant p < 1$. We have

$$\|\mathscr{F}_{\bar{\xi}}^{-1}\psi_{m,l}\mathscr{F}_{\bar{x}}f(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}^p \leqslant \sum_{j\geqslant l-C,\, k\in\mathscr{K}_j^n} \|(\mathscr{F}_{\bar{\xi}}^{-1}\psi_{m,l}\mathscr{F}_{\bar{x}}\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}^p. \tag{4.17}$$

It follows that

$$\begin{aligned}
&\|f(\cdot,0)\|_{M_{p,q}^{s,\alpha}(\mathbb{R}^{n-1})} \\
&\lesssim \left(\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}_l^{n-1}} \langle l\rangle^{\frac{sq}{1-\alpha}} \left(\sum_{j\in\mathbb{Z}_+}\sum_{k\in\mathscr{K}_j^n} \|(\mathscr{F}_{\bar{\xi}}^{-1}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}^p\right)^{q/p}\right)^{1/q} \\
&\lesssim \left(\sum_{j\in\mathbb{Z}_+}\sum_{a=0}^{N_j}\sum_{k\in\mathscr{K}_{j,r_{ja}}^n}\sum_{l\leqslant j+C}\sum_{m\in\mathscr{K}_l^{n-1}} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(\mathscr{F}_{\bar{\xi}}^{-1}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}^q\right)^{1/q}.
\end{aligned} \tag{4.18}$$

For convenience, we write

$$\Upsilon_{ja} := \sum_{k\in\mathscr{K}_{j,r_{ja}}^n}\sum_{l\leqslant j+C}\sum_{m\in\mathscr{K}_l^{n-1}} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(\mathscr{F}_{\bar{\xi}}^{-1}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}^q. \tag{4.19}$$

It follows from Lemma 3.2, the property of $p$-BAPU and (4.7) that

$$\begin{aligned}
\Upsilon_{ja} &\lesssim \sum_{k\in\mathscr{K}_{j,r_{ja}}^n}\ \sum_{a^{1-\alpha}\langle j\rangle^\alpha \lesssim l \lesssim (1+a)^{1-\alpha}\langle j\rangle^\alpha}\ \sum_{m\in\mathscr{K}_l^{n-1}} \langle l\rangle^{\frac{sq}{1-\alpha}} \\
&\qquad \times \langle j\rangle^{q\frac{\alpha(n-1)}{1-\alpha}(\frac{1}{p}-1)} \|(\mathscr{F}_{\bar{\xi}}^{-1}\psi_{m,l})\|_{L^p(\mathbb{R}^{n-1})}^q \|(\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})}^q \\
&\lesssim \sum_{k\in\mathscr{K}_{j,r_{ja}}^n}\ \sum_{a^{1-\alpha}\langle j\rangle^\alpha \lesssim l \lesssim (1+a)^{1-\alpha}\langle j\rangle^\alpha} \langle l\rangle^{\frac{sq}{1-\alpha}} \langle j\rangle^{q\frac{\alpha(n-1)}{1-\alpha}(\frac{1}{p}-1)} \langle l\rangle^{-q\frac{\alpha(n-1)}{1-\alpha}(\frac{1}{p}-1)} \\
&\qquad \times \min\left(\langle l\rangle^{n-2},\ \langle j\rangle^{\frac{\alpha(n-2)}{1-\alpha}} \langle l\rangle^{-\frac{\alpha(n-2)}{1-\alpha}}\right) \langle j\rangle^{\frac{q\alpha}{p(1-\alpha)}} \|\Box_{k,j}^\alpha f\|_{L^p(\mathbb{R}^n)}^q.
\end{aligned} \tag{4.20}$$

If $a \geqslant 1$, then we have

$$\Upsilon_{ja} \lesssim \sum_{k\in\mathscr{K}_{j,r_{ja}}^n} a^{sq-\alpha(n-1)-q\alpha(n-1)(\frac{1}{p}-1)} \langle j\rangle^{\frac{\alpha qs}{1-\alpha}+q\alpha(n-1)(\frac{1}{p}-1)+\alpha(n-1)+\frac{q\alpha}{p(1-\alpha)}} \|\Box_{k,j}^\alpha f\|_{L^p(\mathbb{R}^n)}^q. \tag{4.21}$$

If $a = 0$,

$$\Upsilon_{ja} \lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \sum_{0\leqslant l\lesssim\langle j\rangle^\alpha} \langle l\rangle^{\frac{sq}{1-\alpha}-q\frac{\alpha(n-1)}{1-\alpha}(\frac{1}{p}-1)+(n-2)} \langle j\rangle^{q\frac{\alpha(n-1)}{1-\alpha}(\frac{1}{p}-1)+\frac{q\alpha}{p(1-\alpha)}} \|\Box^\alpha_{k,j} f\|^q_{L^p(\mathbb{R}^n)}. \tag{4.22}$$

Now we divide our discussion into the following four subcases.

*Case 3A.* $qs - \alpha(n-1) - q\alpha(n-1)(1/p-1) \geqslant 0$. If $a \geqslant 1$, we see that the upper bound in (4.21) will be attained at $a \sim \langle j\rangle$. If $a = 0$, the summation on $l$ in (4.22) can be easily controlled. Anyway, we have

$$\Upsilon_{ja} \lesssim \sum_{k\in\mathscr{K}^n_{j,r_{ja}}} \langle j\rangle^{\frac{\alpha q s}{1-\alpha}+\frac{q\alpha}{p(1-\alpha)}} \|\Box^\alpha_{k,j} f\|^q_{L^p(\mathbb{R}^n)}. \tag{4.23}$$

Combining (4.18) and (4.23), we immediately have the result, as desired.

*Case 3B.* $q\alpha s_p - (1-\alpha)(n-1) < qs < q\alpha s_p + \alpha(n-1)$. In this case, using (4.21) and (4.22), we can repeat the procedure as in Case 1B to get the result and we omit the details of the proof.

*Case 3C.* $q\alpha s_p - (1-\alpha)(n-1) = qs$. This case is similar to Case 1C.

*Case 3D.* $q\alpha s_p - (1-\alpha)(n-1) > qs$. We can deal with this case by following the same way as in Case 1D.

*Case 4.* $0 < p < 1$, $q > p$. By Minkowski's inequality, we have

$$\begin{aligned}
&\|f(\cdot,0)\|_{M^{s,\alpha}_{p,q}(\mathbb{R}^{n-1})}\\
&\lesssim \left(\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}^{n-1}_l} \langle l\rangle^{\frac{sq}{1-\alpha}} \left(\sum_{j\in\mathbb{Z}_+}\sum_{k\in\mathscr{K}^n_j} \|(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|^p_{L^p(\mathbb{R}^{n-1})}\right)^{q/p}\right)^{1/q}\\
&\lesssim \left(\sum_{j\in\mathbb{Z}_+}\sum_{k\in\mathscr{K}^n_j} \left(\sum_{l\in\mathbb{Z}_+}\sum_{m\in\mathscr{K}^{n-1}_l} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(\mathscr{F}^{-1}_{\bar{\xi}}\psi_{m,l}) * (\mathscr{F}^{-1}\psi_{k,j}\mathscr{F}f)(\cdot,0)\|^q_{L^p(\mathbb{R}^{n-1})}\right)^{p/q}\right)^{1/p}.
\end{aligned} \tag{4.24}$$

Then we can repeat the procedures as in the proof of Theorem 1.2 and the above techniques in Case 3 to have the result, as desired. The details of the proof are omitted. □

# 5 Proof of Theorem 1.6

Now we prove Theorem 1.6. Now we define the maximum function $M_k^* f$ as follows:

$$M_k^* f = \sup_{y \in \mathbb{Z}^n} \frac{|\Delta_k f(x-y)|}{1+|2^k y|^{n/r}}. \tag{5.1}$$

Taking $y_1 = ... = y_{n-1} = 0$, $y_n = x_n$ in (5.1), we have for $2^{-k-1} \leqslant |x_n| \leqslant 2^{-k}$,

$$|(\Delta_k f)(\bar{x}, 0)| \lesssim |M_k^* f(x)|, \quad \bar{x} = (x_1, \cdots, x_{n-1})$$

Hence

$$\|(\Delta_k f)(\cdot, 0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim \|M_k^* f(\cdot, x_n)\|_{L^p(\mathbb{R}^{n-1})}, \tag{5.2}$$

Integrating (5.2), one has that

$$\|(\Delta_k f)(\cdot, 0)\|^p_{L^p(\mathbb{R}^{n-1})} \lesssim 2^k \int_{\mathbb{R}} \|M_{k,t}^* f(\cdot, x_n)\|^p_{L^p(\mathbb{R}^{n-1})} dx_n,$$

Hence

$$\|(\Delta_k f)(\cdot, 0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim 2^{k/p} \|M_k^* f\|_{L^p(\mathbb{R}^n)}. \tag{5.3}$$

Write $\varphi'_k(\bar{x})$ as the BAPU functions in $\mathbb{R}^{n-1}$. Then for fixed $k$, we have

$$\begin{aligned}(\mathscr{F}_{\bar{\xi}}^{-1} \varphi'_k \mathscr{F}_{\bar{x}})(\bar{x}, 0) &= \sum_{l=k-1}^{\infty} (\mathscr{F}_{\bar{\xi}}^{-1} \varphi'_k \mathscr{F}_{\bar{x}} \mathscr{F}^{-1} \varphi_l \mathscr{F})(\bar{x}, 0) \\ &= \sum_{l=k-1}^{\infty} (\mathscr{F}_{\bar{\xi}}^{-1} \varphi'_k) * (\mathscr{F}^{-1} \varphi_l \mathscr{F})(\bar{x}, 0)\end{aligned}$$

*Case 1.* $1 \leqslant p \leqslant \infty$. Using Young's inequality, (3.1) and (5.3), we obtain

$$\begin{aligned}&\|\mathscr{F}_{\bar{\xi}}^{-1} \varphi'_k \mathscr{F}_{\bar{x}} f(\bar{x}, 0)\|_{L^p(\mathbb{R}^{n-1})} \\ &\lesssim \sum_{l=k-1}^{\infty} \|\mathscr{F}_{\bar{\xi}}^{-1} \varphi'_k\|_{L^1(\mathbb{R}^{n-1})} \|\mathscr{F}^{-1} \varphi_l \mathscr{F} f\|_{L^p(\mathbb{R}^{n-1})} \\ &\lesssim \sum_{l=k-1}^{\infty} \|M_l^* f\|_{L^p(\mathbb{R}^n)} \\ &\lesssim \sum_{l=k-1}^{\infty} 2^{l/p} \|\Delta_l f\|_{L^p(\mathbb{R}^n)}.\end{aligned}$$

Hence,

$$\|f(\bar{x}, 0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left( \sum_{k=0}^{\infty} 2^{skq} \left( \sum_{l=k-1}^{\infty} 2^{l/p} \|\Delta_l f\|_{L^p(\mathbb{R}^n)} \right)^q \right)^{1/q}.$$

If $0 < q \leqslant 1$, then

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left( \sum_{l=-1}^{\infty} \sum_{k=0}^{l+1} 2^{skq} 2^{lq/p} \|\Delta_l f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q}.$$

If $s = 0$, then

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left( \sum_{l=-1}^{\infty} l 2^{lq/p} \|\Delta_l f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q} \lesssim \|f\|_{\tilde{B}^{1/p}_{p,q}(\mathbb{R}^n)}.$$

In the case $s < 0$,

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left( \sum_{l=-1}^{\infty} 2^{lq/p} \|\Delta_l f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q} \lesssim \|f\|_{B^{1/p}_{p,q}(\mathbb{R}^n)}.$$

If $1 \leqslant q \leqslant \infty$, using Minkowski's inequality,

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \sum_{l=-1}^{\infty} \left( \sum_{k=0}^{l+1} 2^{skq} 2^{lq/p} \|\Delta_l f\|^q_{L^p(\mathbb{R}^n)} \right)^{1/q}$$

then

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \begin{cases} \|f\|_{\tilde{B}^{1/p}_{p,1}(\mathbb{R}^n)} & s = 0, \\ \|f\|_{B^{1/p}_{p,1}(\mathbb{R}^n)} & s < 0. \end{cases}$$

*Case 2.* $0 < p < 1$.

$$\begin{aligned} &\|\mathscr{F}^{-1}_{\bar{\xi}} \varphi'_k \mathscr{F}_{\bar{x}} f(\bar{x},0)\|^p_{L^p(\mathbb{R}^{n-1})} \\ &\lesssim \sum_{l=k-1}^{\infty} 2^{l(n-1)(1/p-1)p} \|\mathscr{F}^{-1}_{\bar{\xi}} \varphi'_k\|^p_{L^p(\mathbb{R}^{n-1})} \|\mathscr{F}^{-1} \varphi_l \mathscr{F} f\|^p_{L^p(\mathbb{R}^{n-1})} \\ &\lesssim \sum_{l=k-1}^{\infty} 2^{l(n-1)(1-p)} 2^{k(n-1)(p-1)} 2^l \|M^*_l f\|^p_{L^p(\mathbb{R}^n)} \\ &\lesssim \sum_{l=k-1}^{\infty} 2^{l(n-1)(1-p)} 2^{k(n-1)(p-1)} 2^l \|\Delta_l f\|^p_{L^p(\mathbb{R}^n)}. \end{aligned}$$

It follows that

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left( \sum_{k=0}^{\infty} 2^{skq} \left( \sum_{l=k-1}^{\infty} 2^{l(n-1)(1-p)} 2^{k(n-1)(p-1)} 2^l \|\Delta_l f\|^p_{L^p(\mathbb{R}^n)} \right)^{q/p} \right)^{1/q}.$$

If $q \leqslant p$, one has that

$$\begin{aligned}
&\|f(\bar{x},0)\|^q_{B^s_{p,q}(\mathbb{R}^{n-1})}\\
&\lesssim \sum_{k=0}^{\infty}\sum_{l=k-1}^{\infty} 2^{l(n-1)(1/p-1)q}2^{k(s+(n-1)(1-1/p))q}2^{lq/p}\|\Delta_l f\|^q_{L^p(\mathbb{R}^n)}\\
&\lesssim \sum_{l=-1}^{\infty}\sum_{k=0}^{l+1} 2^{l(n-1)(1/p-1)q}2^{k(s+(n-1)(1-1/p))q}2^{lq/p}\|\Delta_l f\|^q_{L^p(\mathbb{R}^n)}.
\end{aligned}$$

Recall that we write $s_p = (n-1)(1/(p\wedge 1)-1)$, then

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \begin{cases} \|f\|_{\tilde{B}^{s_p+1/p}_{p,q}(\mathbb{R}^n)} & s = s_p,\\ \|f\|_{B^{s_p+1/p}_{p,q}(\mathbb{R}^n)}, & s < s_p. \end{cases}$$

If $q \geqslant p$, using Minkowski's inequality, we have

$$\begin{aligned}
&\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})}\\
&\lesssim \left[\sum_{l=-1}^{\infty}\left(\sum_{k=0}^{\infty}\left(\chi(k\leqslant l)2^{l(n-1)(1-p)}2^{k(s+(n-1)(p-1))}2^l\|\Delta_l f\|^p_{L^p(\mathbb{R}^n)}\right)^{q/p}\right)^{p/q}\right]^{1/p}\\
&\lesssim \left[\sum_{l=-1}^{\infty}\sum_{k=0}^{l+1} 2^{l(n-1)(1/p-1)p}2^{k(s+(n-1)(1-1/p))p}2^l\|\Delta_l f\|^q_{L^p(\mathbb{R}^n)}\right]^{1/p}.
\end{aligned}$$

Therefore,

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \begin{cases} \|f\|_{\tilde{B}^{s_p+1/p}_{p,p}(\mathbb{R}^n)}, & s = s_p,\\ \|f\|_{B^{s_p+1/p}_{p,p}(\mathbb{R}^n)}, & s < s_p. \end{cases}$$

In the case $1 < p < \infty$, we define the maximum function $M^*_{k,t}f$ as follows:

$$M^*_{k,t}f = \sup_{y\in\mathbb{Z}^n}\frac{|\Delta_{k,t}f(x-y)|}{1+|2^k y|^{n/r}}. \tag{5.4}$$

Taking $y_1 = ... = y_{n-1} = 0$, $y_n = x_n$ in (5.4), we have for $2^{-k-1} \leqslant |x_n| \leqslant 2^{-k}$,

$$|(\Delta_{k,t}f)(\bar{x},0)| \lesssim |M^*_{k,t}f(x)|, \quad \bar{x} = (x_1,\cdots,x_{n-1})$$

Hence

$$\|(\Delta_{k,t}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim \|M^*_{k,t}f(\cdot,x_n)\|_{L^p(\mathbb{R}^{n-1})}, \tag{5.5}$$

Integrating (5.5), one has that

$$\|(\Delta_{k,t}f)(\cdot,0)\|^p_{L^p(\mathbb{R}^{n-1})} \lesssim 2^k\int_{\mathbb{R}}\|M^*_{k,t}f(\cdot,x_n)\|^p_{L^p(\mathbb{R}^{n-1})}dx_n,$$

Hence

$$\|(\Delta_{k,t}f)(\cdot,0)\|_{L^p(\mathbb{R}^{n-1})} \lesssim 2^{k/p}\|M^*_{k,t}f\|_{L^p(\mathbb{R}^n)}. \tag{5.6}$$

Let $\chi'_{k,t}(\bar{x})$ as the characteristic functions in $\mathbb{R}^{n-1}$. Then for fixed $k$ and $t$, we have

$$\begin{aligned}(\mathscr{F}_{\bar{\xi}}^{-1}\chi'_{k,t}\mathscr{F}_{\bar{x}})(\bar{x},0) &= \sum_{l=k}^{\infty}(\mathscr{F}_{\bar{\xi}}^{-1}\chi'_{k,t}\mathscr{F}_{\bar{x}}\mathscr{F}^{-1}\chi_{l,t}\mathscr{F})(\bar{x},0)\\ &= \sum_{l=k}^{\infty}(\mathscr{F}_{\bar{\xi}}^{-1}\chi'_{k,t}) * (\mathscr{F}^{-1}\chi_{l,t}\mathscr{F})(\bar{x},0)\end{aligned}$$

Using Young's inequality, (3.1) and (5.6), we obtain

$$\begin{aligned}&\|\mathscr{F}_{\bar{\xi}}^{-1}\chi'_{k,t}\mathscr{F}_{\bar{x}}f(\bar{x},0)\|_{L^p(\mathbb{R}^{n-1})}\\ &\lesssim \sum_{l=k}^{\infty}\|\mathscr{F}_{\bar{\xi}}^{-1}\chi'_{k,t}\|_{L^1(\mathbb{R}^{n-1})}\|\mathscr{F}^{-1}\chi_{l,t}\mathscr{F}f\|_{L^p(\mathbb{R}^{n-1})}\\ &\lesssim \sum_{l=k}^{\infty}\|M^*_{l,t}f\|_{L^p(\mathbb{R}^n)}\\ &\lesssim \sum_{l=k}^{\infty}2^{l/p}\|\Delta_{l,t}f\|_{L^p(\mathbb{R}^n)}.\end{aligned}$$

Hence,

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left(\sum_{k=0}^{\infty}\sum_{t=1}^{T}2^{skq}\left(\sum_{l=k}^{\infty}2^{l/p}\|\Delta_{l,t}f\|_{L^p(\mathbb{R}^n)}\right)^q\right)^{1/q}.$$

If $0 < q \leqslant 1$, then

$$\begin{aligned}&\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})}\\ &\lesssim \left(\sum_{l=0}^{\infty}\sum_{k=0}^{l-1}\sum_{t=1}^{2^n}2^{skq}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)} + \sum_{l=0}^{\infty}\sum_{k=l}\sum_{t=2^n+1}^{T}2^{skq}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}.\end{aligned}$$

If $s = 0$, then

$$\begin{aligned}\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} &\lesssim \left(\sum_{l=0}^{\infty}\sum_{t=1}^{2^n}l2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)} + \sum_{l=0}^{\infty}\sum_{t=2^n+1}^{T}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}\\ &\lesssim \|f\|_{\tilde{B}^{1/p,1/p}_{p,q}}.\end{aligned}$$

In the case $s < 0$,

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \left(\sum_{l=0}^{\infty}\sum_{t=1}^{2^n}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)} + \sum_{l=0}^{\infty}\sum_{t=2^n+1}^{T}2^{slq}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}$$

$$\lesssim \|f\|_{B^{1/p,s+1/p}_{p,q}}.$$

If $1 \leqslant q \leqslant \infty$, using Minkowski's inequality,

$$\begin{aligned}
&\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \\
&\lesssim \sum_{l=0}^{\infty}\left(\sum_{k=0}^{l}\sum_{t=1}^{T} 2^{skq}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q} \\
&\lesssim \sum_{l=0}^{\infty}\left(\sum_{k=0}^{l-1}\sum_{t=1}^{2^n} 2^{skq}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)} + \sum_{k=l}\sum_{t=2^n+1}^{T} 2^{skq}2^{lq/p}\|\Delta_{l,t}f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}.
\end{aligned}$$

then

$$\|f(\bar{x},0)\|_{B^s_{p,q}(\mathbb{R}^{n-1})} \lesssim \begin{cases} \|f\|_{\tilde{B}^{1/p,1/p}_{p,1}(\mathbb{R}^n)} & s=0, \\ \|f\|_{B^{1/p,s+1/p}_{p,1}(\mathbb{R}^n)} & s<0. \end{cases}$$

□

**Acknowledgment.** This work is partially supported by the Marie Curie Excellence Project EUCETIFA, MEXT-CT-2004-517154 of the European Commission. The second and third named authors are supported in part by the National Science Foundation of China, grant 10571004; and the 973 Project Foundation of China, grant 2006CB805902.

# References

[1] L. Borup, M. Nielsen, Banach frames for multivariante $\alpha$-modulation spaces, J. Math. Anal. Appl., **321** (2006), 880-895.

[2] J. Bergh and J. Löfström, Interpolation Spaces, Springer–Verlag, 1976.

[3] H. G. Feichtinger and P. Gröbner, Banach spaces of distributions defined by decomposition methods. I, Math. Nachr. 132 (1985) 207-237.

[4] H. G. Feichtinger, Banach spaces of distributions defined by decomposition methods. II, Math. Nachr. 123 (1987) 97-120.

[5] H. G. Feichtinger, Modulation spaces on locally compact Abelian group, Technical Report, University of Vienna, 1983. Published in: "Proc. Internat. Conf. on Wavelet and Applications", 99–140. New Delhi Allied Publishers, India, 2003. http://www.unive.ac.at/nuhag-php/bibtex/ open_files/fe03-1_modspa03.pdf.

[6] H. G. Feichtinger, Banach spaces of distributions of Wiener's type and interpolation, Proceedings on Functional Analysis and Approximation (Oberwilfach, 1980), 153–165. Internat. Ser. Numer. Math., 60 Birkäuser, Basel-Boston, Mass., 1981.

[7] Massimo Fornasier, Banach frames for $\alpha$–modulation spaces, Appl. Comput. Harmon. Anal., Vol.22 No.2, (2007) p.157–175.

[8] M. Frazier and B. Jawerth, Decomposition of Besov spaces, Indiana Univ. Math. J. 34 (1985), 777-799.

[9] Yevgeniy V. Galperin and Salti Samarah, Time-frequency analysis on modulation spaces $M_m^{p,q}$, Appl. Comput. Harmon. Anal., Vol.16 No.1, (2004) p.1–18.

[10] P. Gröbner, Banachräume Glatter Funktionen und Zerlegungsmethoden, Doctoral thesis, University of Vienna, 1992.

[11] M. Kobayashi, Modulation spaces $M^{p,q}$ for $0 < p, q \leqslant \infty$, J. of Funct. Spaces and Appl. **4** (2006) no.3, 329–341.

[12] L. Päivärinta and E. Somersalo, A generalization of the Calderön-Vaillancourt theorem to $L^p$ and $h^p$, Math.Nachr. 138 (1988) 145-156.

[13] H. Rauhut, Coorbit Space Theory for Quasi-Banach Spaces, Studia Math., Vol.180, No.3, (2007) p.237-253.

[14] H. Triebel, Theory of Function Spaces, Birkhäuser–Verlag, 1983.

[15] H. Triebel, Complex interpolation and Fourier multiplier for the spaces $B_{p,q}^s$ and $F_{p,q}^s$ of Besov–Hardy–Sobolev Type: The case $0 < p, q \leqslant \infty$, Math. Z., **176** (1981), 495–510.

[16] H. Triebel, Modulation spaces on the Euclidean $n$-spaces, Z. Analysis Anwendungen, **2** (1983), 443–457.

[17] Baoxiang Wang, Lifeng Zhao and Boling Guo, Isometric decomposition operators, function spaces $E_{p,q}^{\lambda}$ and applications to nonlinear evolution equations, J. Funct. Anal., **233** (2006), 1–39.

[18] Baoxiang Wang and Henryk Hudzik, The global Cauchy problem for the NLS and NLKG with small rough data, J. Differential Equations, **231** (2007), 36–73.

[19] Baoxiang Wang and Chunyan Huang, Frequency-uniform decomposition method for the generalized BO, KdV and NLS equations J. Differential Equations, **231** (2007), 36–73.